

**Exercise 3—Due in section from 9 March**

Name: \_\_\_\_\_

**The Winter Skies**

Section: \_\_\_\_\_

**Step 1.** Calibrate your hand.

In this exercise, you will estimate the altitude of stars above the horizon, using your hand to measure angles. So the first thing to do is to figure out how many degrees across your hand is. Stand directly between two objects (the two sides of a wide door, two trees, two goalposts, etc.), so that they are  $180^\circ$  apart. Make a fist with your hand, with your thumb covering the top of your fist, and extend it to arm's length. Then sight down your arm with one eye. To determine how many degrees of angle your hand takes up, count how many hand widths it takes to cover  $180^\circ$  of angle. Keep moving your hand one width to the left or right at a time, using background objects as reference points to keep track.

$$\text{_____ handwidths} = H = 180^\circ.$$

The quantity  $180^\circ/H$  = the angular width of your hand = \_\_\_\_\_.

It's OK to round to the nearest degree. The answer will be about  $10^\circ$  for most people.

**Step 2.** Study a star chart.

Your textbook has a nice set of starcharts in Appendix J. From the warm cozy comfort of your own room, have a look at the winter chart and try to get a feel for where the different constellations will be in the sky before you actually step out into the cold night air. If a star is in the center of the circle (like Capella), that means that it will be close to the zenith if you observe at 8 pm (i.e. its altitude is  $90^\circ$ ). A star halfway from the center to one side of the circle is  $45^\circ$  above that direction on the horizon.

Most likely, the stars will be shifted a little from their positions in the star charts because they appear to rotate about the north celestial pole each night. When using the combinations of dates and times at the bottom of the charts, notice that a shift of one month equals two hours. If you are crazy enough to be observing before dawn, you'll need to use the spring chart.

**Step 3.** Measure the altitude and azimuth of several bright stars and planets.

Your task is to measure the altitude and azimuth of the objects listed in the table on the next page. To do this, you will need to find a dark location with a good view of the horizon. Don't forget to dress **warmly** (i.e. really warmly). It's a good idea to bring a flashlight and your starcharts, since it takes a little practice to find your way around the night sky.

As many of you are well aware, Ithaca is not famous for its clear skies, especially in the winter, so don't pass up the first good night you get. It might be the only clear night before the assignment is due.

Record the date and time of your observation:

Date: \_\_\_\_\_

Time: \_\_\_\_\_

For each object in the table below, record its altitude and azimuth. The definitions below might help you. As a bonus, see if you can detect a color difference between Betelgeuse and Rigel, and between Castor and Pollux. The values for Polaris are already filled in.

**Altitude** — the angular distance of an object above the horizon directly beneath it. Record this measurement in degrees from the horizon to the object. Realistically, you will only be able to estimate the altitude to the nearest five degrees.

**Azimuth** — the direction of the horizon directly beneath an object. Record your answers as "NE", "E", "SE", etc. If an object is at the zenith (altitude=90°), then it doesn't really have an azimuth.

Target	Altitude	Azimuth	Color
Polaris ( $\alpha$ UMi)	<u>42°</u>	<u>N</u>	
Capella ( $\alpha$ Aur)	_____	_____	
Aldebaran ( $\alpha$ Tau)	_____	_____	
Betelgeuse ( $\alpha$ Ori)	_____	_____	_____
Rigel ( $\beta$ Ori)	_____	_____	_____
Sirius ( $\alpha$ CMa)	_____	_____	
Pollux ( $\alpha$ Gem)	_____	_____	_____
Castor ( $\beta$ Gem)	_____	_____	_____
Regulus ( $\alpha$ Leo)	_____	_____	
Venus	_____	_____	
Saturn	_____	_____	

We're lucky to have a comet in the sky right now. Comet Lulin will be moving through Libra in early February, then Virgo and Leo later in the month. On 27 February, it will be within a degree of Regulus. Right now, it's a little fuzz ball easily visible with binoculars. Soon, on a good night, it should be a naked-eye object by mid February. Can you find it?